

On the odderon intercept in the variational approach

M.A.Braun

Department of High Energy Physics, University of St. Petersburg,
198904 St. Petersburg, Russia.

January 1998

Abstract.

The odderon intercept α is recalculated by the variational method based on conformal invariance. The final result is $\alpha = 1 - 0.339 \alpha_s N / \pi$ in contrast to the published value above unity.

SPbU-IP- 1998/3

1 Introduction

As shown in [1,2], at least in the high-colour limit, all states formed by any number of reggeized gluons reduce to two basic ones: the pomeron for the positive signature and the odderon for the negative signature. The intercepts of these two states are fundamental for the high-energy behavior of cross-sections in the perturbative QCD. Unlike the pomeron intercept, known since more than two decades ago, the odderon intercept has not been calculated exactly until now, despite much activity around this problem and most sophisticated mathematical tools used [3-6]. The only published results refer to estimates made in the variational approach [7-9]. In paper [8], based on the conformal property of the odderon [7], the obtained intercept lies above unity. In the following we present it in the form

$$\alpha = 1 - \frac{\alpha_s N}{\pi} \frac{3}{2} \epsilon \quad (1)$$

Here α_s is the strong coupling constant, N is the number of colours; ϵ has a meaning of the odderon "energy" per gluon in units $\alpha_s N/\pi$. The corresponding energy for the pomeron is $-4 \ln 2$. It is negative, which implies the intercept above unity and cross-sections growing with energy like s^Δ where $\Delta = -(\alpha_s N/\pi)\epsilon$. For the odderon in [8] it was obtained

$$\epsilon = -0.25 \quad (2)$$

which is also negative but considerably greater than the pomeron energy. If correct, it would correspond to the negative signature cross-sections also rising with energy, although much slower than the positive signature ones.

In the subsequent paper [9] the conformal properties of the odderon were not taken into account. Instead its wave function was presented as a sum of one-gluon "orbitals", which made it possible to take a very large number of variational parameters. A positive value was obtained for the energy:

$$\epsilon = 0.298 \quad (3)$$

In principle, due to the variational nature of the calculations, this value does not contradict (2). However in [9] certain theoretical arguments were also presented in favour of the odderon intercept being smaller than unity.

Guided by the evident conflict of these theoretical arguments with the result (2), we undertook to repeat the variational calculations in the conformal approach of [7,8], taking both the same trial function and the ones with more parameters. It turned out that the result (2) is in error. Correct calculations with the same trial function (with 2 parameters) give

$$\epsilon = 0.2286 \pm 0.0001 \quad (4)$$

with an opposite sign as compared to (2). Inclusion of more parameters (up to 8) allows to slightly lower this value down to

$$\epsilon = 0.2262 \pm 0.0001 \quad (5)$$

The uncertainties in (4) and (5) refer to errors in numerical calculations. In view of a very small change in ϵ upon introducing more parameters, one is inclined to take (5) as a final value for the odderon energy with a precision of 0.1%.

The study of the odderon energy in the conformal approach, in fact, presents a formidable calculational problem, deserving some attention. In this note we briefly discuss the difficulties involved and some intermediate steps in obtaining our results.

2 Basic equations

Basic equations for the odderon energy in the conformal approach were derived in [7]. We reproduce them here to fix our notations. The odderon energy is to be sought as a minimum of the energy functional

$$E = \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} d\nu \epsilon_n(\nu) |\alpha_n(\nu)|^2 \quad (6)$$

Here

$$\alpha_n(\nu) = \int_0^{\infty} dr r^{-2-2i\nu} \int_0^{2\pi} d\phi e^{-in\phi} \left(i\nu + \frac{n+1}{2} + re^{i\phi} \left(h - i\nu - \frac{n-1}{2} \right) \right) \left(i\nu - \frac{n-1}{2} \right) (-\tilde{h} + i\nu - \frac{n-1}{2}) Z(r, \phi) \quad (7)$$

Z is the odderon wave function and

$$\epsilon_n(\nu) = 2 \operatorname{Re} \left(\psi \left(\frac{1+|n|}{2} + i\nu \right) - \psi(1) \right) \quad (8)$$

The normalization condition is

$$1 = D = \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} d\nu |\alpha_n(\nu)|^2 \quad (9)$$

In these formulas h and $\tilde{h}$ are the two conformal weights, which are taken to be equal to $1/2$. The trial function Z then takes the form

$$Z(r, \phi) = r^{1/3} r_1^{1/3} z(r, \phi) \quad (10)$$

where

$$r_1^2 = 1 + r^2 - 2r \cos \phi \quad (11)$$

and $z(r, \phi) = z(r, r^*)$ should be invariant under the transformations of complex r : $r \rightarrow -1/r$ and $r \rightarrow 1 - r$. This can be achieved by taking $z = z(a)$ where

$$a = \frac{r^2 r_1^2}{(1 + r^2)(1 + r_1^2)(r^2 + r_1^2)} \quad (12)$$

We take the trial function $z(a)$ in the form

$$z(a) = \sum_{k=1}^{N-1} c_k a^{k/2-1/6} + c_N a^{5/6} \ln a \quad (13)$$

with $N-1$ variational parameters c_k (one of c_k is determined by the normalization condition). For $N = 3$ this function coincides with the one used in [8].

The basic quantity α given by Eq. (7) can be presented in the form

$$\alpha_n(\nu) = \left(\frac{n}{2} - i\nu\right) \left(\frac{n-1}{2} - i\nu\right) \left[\left(\frac{n+1}{2} + i\nu\right) f_n^{(1)}(\nu) + \left(1 - \frac{n}{2} - i\nu\right) f_{n-1}^{(0)}(\nu)\right] \quad (14)$$

where

$$f_n^{(k)} = \int_0^\infty dr r^{-1-k-2i\nu} \int_0^{2\pi} d\phi e^{-in\phi} Z(r, \phi) \quad (15)$$

Function $f_n^{(k)}(\nu)$ has the following properties. Since Z is real, $(f_n^{(k)}(\nu))^* = f_{-n}^{(k)}(-\nu)$ and since Z only depends on ϕ through $\cos \phi$, we have $f_n^{(k)}(\nu) = f_{-n}^{(k)}(\nu)$. Under $r \rightarrow 1/r$ the argument a stays invariant, from which it follows that $Z \rightarrow Z/r$. As a result we obtain $f_n^{(1)}(\nu) = f_n^{(0)}(-\nu) = (f_n^{(0)})^*$. Using these properties we can restrict the summation over n and integration over ν to nonnegative values. The appearing coefficient 2 cancels in dividing E by D so that we need not take it into account. The value of the $|\alpha_n(\nu)|^2$ can evidently be expressed via a single function $f_n^{(0)}(\nu)$, which will simply be denoted as $f_n(\nu)$ in the following. In this manner we obtain our basic formulas for E and D :

$$E = \sum_{n=0}^{\infty} \int_0^\infty d\nu \epsilon_n(\nu) p_n(\nu) \quad (16)$$

where for $n > 0$

$$p_n(\nu) = \left(\frac{n^2}{4} + \nu^2\right) \left(\frac{(n-1)^2}{4} + \nu^2\right) \left[\left(\frac{(n+1)^2}{4} + \nu^2\right) |f_n(\nu)|^2 + \left(\frac{(n-2)^2}{4} + \nu^2\right) |f_{n-1}(\nu)|^2 + 2 \operatorname{Re} \left(\frac{n+1}{2} - i\nu\right) \left(\frac{2-n}{2} - i\nu\right) f_n(\nu) f_{n-1}(\nu)\right] \quad (17)$$

and

$$p_0(\nu) = \nu^2 \left(\frac{1}{4} + \nu^2\right) \left[\left(\frac{1}{4} + \nu^2\right) |f_0(\nu)|^2 + (1 + \nu^2) |f_1(\nu)|^2 + 2 \operatorname{Re} \left(\frac{1}{2} - i\nu\right) (1 - i\nu) f_0(\nu) f_1(\nu)\right] \quad (18)$$

The normalization functional D has the same form (16) with $\epsilon_n(\nu) \rightarrow 1$. Thus calculation of the odderon energy requires calculation of functions $f_n(\nu)$ and $\epsilon_n(\nu)$.

The main technical difficulty is a double Fourier transform (15). The energy $\epsilon_n(\nu)$ in E , Eq. (16), monotonously grows both with n and ν . It is negative only for $n = 0$ and small enough values of ν . So the problem with this formalism is that cutting in (16) summation over n and integration over ν by some maximal values n_m and ν_m , one always gets smaller E than the exact value, corresponding to n_m and $\nu_m \rightarrow \infty$. Therefore in the course of the

calculation one always approaches to the variational value of ϵ *from below*. As we shall see in the following, in fact, rather high values of n_m and ν_m are necessary to obtain ϵ with some degree of accuracy ¹. On the other hand, with high n and ν , the double Fourier transform (15) becomes very difficult, especially having in mind that, due to the factors in (17), two first terms in the asymptotic expansion of α at high n and ν cancel. As a result, as mentioned in the Introduction, a trustworthy calculation of E and D turns out to be very complicated, in spite of its superficial transparency.

The crucial point in obtaining reasonable results has been using analytic asymptotic expansions for f at high n and ν , which are discussed in the next section.

3 Asymptotics at large n and ν

Passing to variable $\rho = -\ln r$ and introducing 2-dimensional vectors $x = (\rho, \phi)$. and $w = (z, n) = (2\nu, n)$ we rewrite (15) as

$$f_n(\nu) = \int_{-\infty}^{+\infty} d\rho \int_{-\pi}^{+\pi} d\phi e^{iwx} Z(x) \quad (19)$$

Evidently for the asymptotics at high n and ν the integration point $x = 0$ is essential.

At $x \rightarrow 0$, keeping terms up to third order in small ρ and ϕ , we have

$$\begin{aligned} r^{1/3} &= 1 - (1/3)\rho + (1/18)\rho^2 - (1/162)\rho^3 \\ r_1^{1/3} &= x^{1/3}(1 - (1/6)\rho + (1/36)\rho^2 - (1/72)\phi^2 - (1/324)\rho^3 + (1/432)\rho\phi^2) \\ a^p &= (1/2)^p x^{2p}(1 - p(29/12)\rho^2 - p(25/12)\phi^2) \end{aligned}$$

(no terms of the third order appear in a). In these formulas $x = \sqrt{\rho^2 + \phi^2}$. As a result, at small x

$$Z_p = (rr_1)^{1/3} a^p = (1/2)^p x^{2p+1/3} (1 - (1/2)\rho + a\rho^2 + b\phi^2 + c\rho^3 + d\rho\phi^2) \quad (20)$$

where

$$a = 5/36 - (29/12)p; \quad b = -1/72 - (25/12)p; \quad c = -1/36 + (29/24)p; \quad d = 1/144 + (25/24)p$$

For the term with a logarithm, in the same manner we obtain

$$\begin{aligned} \tilde{Z}_p &= (rr_1)^{1/3} a^p \ln a = (1/2)^p x^{2p+1/3} [\ln(x^2)(1 - (1/2)\rho + a\rho^2 + b\phi^2 + c\rho^3 + d\rho\phi^2) + \\ &\quad \ln(1/2) - (1/2) \ln(1/2) \rho + a_1\rho^2 + b_1\phi^2 + c_1\rho^3 + d_1\rho\phi^2] \end{aligned} \quad (21)$$

¹This is the reason why in [7] the negative value (2) was obtained. The authors chose too small values of n_m and ν_m

where

$$f_1 = f \ln(1/2) + \delta_f, \quad f = a, b, c, d$$

and

$$\delta_a = -29/12, \quad \delta_b = -25/12, \quad \delta_c = 29/24, \quad \delta_d = 25/24$$

Inserting these expressions into the integral (19) and extending the integration over ϕ to the whole real axis one obtains the asymptotical expansion of different terms in $f_n(\nu)$. In particular the asymptotical expansion of the term originating from Z_p is found as

$$f_n^{(p)}(\nu) = c_p(1 + (1/2)id/dz - ad^2/dz^2 - bd^2/dn^2 + icd^3/dz^3 + idd^3/dzdn^2)w^{-\alpha-1}$$

where $\alpha = 2p + 4/3$ and

$$c_p = 2^{\alpha+1-p} \Gamma^2(1/2 + \alpha/2) \cos(\pi\alpha/2)$$

Doing the derivatives, one obtains finally

$$\text{Re} f_n^{(p)}(\nu) = c_p w^{-\alpha-1} \left(1 + (\alpha+1)w^{-4} [z^2(b - a(\alpha+2) + n^2(a - b(\alpha+2)))] \right) \quad (22)$$

$$\text{Im} f_n^{(p)}(\nu) = -(1/2)c_p(\alpha+1)zw^{-\alpha-3} \left(1 + 2(\alpha+3)w^{-4} [z^2(c(\alpha+2) - d) + n^2(d(\alpha+4) - 3c)] \right) \quad (23)$$

For the term with a logarithm at $p = 5/6$ only the part with $\ln x^2$ contributes. The result coincides with the formula above with $p = 5/6$ and the constant c_p substituted by $16\pi(1/2)^{5/6}$.

If one puts these asymptotic expressions into (17) one finds that the two leading terms coming from $Z_{1/3}$ cancel. Numerically the asymptotic expansion begins to work at rather high values of n and ν : $\sqrt{n^2 + 4\nu^2} > 30$.

4 Numerical procedure

According to (15) function $f_n(\nu)$ is defined as an integral

$$f_n = \int_0^\infty dr r^{-1-2i\nu} \int_0^{2\pi} d\phi e^{-in\phi} Z(r, \phi) \quad (24)$$

The double Fourier transform (24) was done numerically in the interval $0 \leq n < 30$, $0 < \nu < 15$.

At $r > 1$ we transform the integration variable $r \rightarrow 1/r$, under which $Z \rightarrow Z/r$. Then the integration over r becomes restricted to the interval $0 < r < 1$. Using properties of Z we can also integrate only over $0 < \phi < \pi$ and substitute $e^{-in\phi}$ by $2 \cos n\phi$. We finally use that Z linearly goes to zero as $r \rightarrow 0$ and introduce a new function

$$\zeta(r, \phi) = Z(r, \phi)/r \quad (25)$$

In terms of ζ we then have

$$f_n = 2 \int_0^1 dr (r^{-2i\nu} + r^{-1+2i\nu}) \int_0^\pi d\phi \cos in\phi \zeta(r, \phi) \quad (26)$$

The integrand is evidently singular at $r = 0$. To soften this behaviour we use that at small r

$$\zeta(r, \phi) = \zeta_0 - \zeta_1 r \ln r + \zeta_2 r + \mathcal{O}(r^2 \ln r) \quad (27)$$

where

$$\zeta_0 = c_1(1/2)^{1/3}, \quad \zeta_1 = -2c_N(1/2)^{5/6}, \quad \zeta_2 = (1/2)^{5/6}(c_2 - c_N \ln 2) \quad (28)$$

In (26) we subtract from ζ the first term of expansion (27) in the part of the integrand with $r^{-2i\nu}$ and the first three terms of this expansion in the part with $r^{-1+2i\nu}$, adding to $f_n(\nu)$ the result of the integration of subtracted terms, which can be found explicitly. In this way we come to our final formula

$$\begin{aligned} f_n(\nu) = & 2 \int_0^\pi d\phi \cos in\phi \int_0^1 dr (r^{-2i\nu} (\zeta(r, \phi) - \zeta_0) \\ & + r^{-1+2i\nu} (\zeta(r, \phi) - \zeta_0 + \zeta_1 r \ln r - \zeta_2 r)) + \\ & \delta_{n0} 2\pi \left(\frac{\zeta_0}{2i\nu(1-2i\nu)} + \frac{\zeta_1}{(1+2i\nu)^2} + \frac{\zeta_2}{1+2i\nu} \right) \end{aligned} \quad (29)$$

Eq. (29) was used for numerical calculation of $f_n(\nu)$ in the above mentioned interval of n and ν . Integrations were performed by dividing the rectangle $0 < r < 1$, $0 < \phi < \pi$ into an $M \times M$ grid, interpolating ζ quadratically on the grid and then doing the integrals explicitly. The maximal value of M was 640. The achieved accuracy was about 10^{-5} .

Thus calculated values of $f_n(\nu)$ were summed over n and integrated over ν as indicated in (16)-(18) to obtain E and D . Stable results were obtained with the quite high maximal values $n_m = 300$ and $\nu_m = 150$. In the part of n, ν space outside the rectangle $0 \leq n < 30$, $0 < \nu < 15$, the asymptotic expressions (22) and (23) were used for $f_n(\nu)$.

As a result we calculated E and D as a quadratic form in the variational parameters c_k . Afterwards the minimal value ϵ of E subject to condition $D = 1$ was found by standard methods.

The results for different number of parameters N are presented in the Table. To see the importance of the high n, ν region, we also present the values of energy ϵ_1 calculated without the asymptotical region, that is restricting to $0 \leq n < 30$, $0 < \nu < 15$. The standard precision corresponds to the r, ϕ grid 320×320 . To clarify the accuracy achieved we also present the results with a double precision (the grid 640×640) for $N = 5$.

Inspecting these results we see that the final accuracy in energy is of the order $7 \cdot 10^{-5}$. This implies that taking $N > 6$ has no sense within the precision achieved, since the corresponding change in energy is of the same order or less.

So our conclusion is that the variational odderon energy with a trial function (13) is given by (5) and that with the accuracy achieved in the course of numerical integration, as described above, the maximal number of terms to be taken in the trial function is $N = 6$, although already with $N = 3$ used in [8] the energy is obtained up to 1%.

5 Acknowledgements

The author expresses his deep gratitude to P.Gauron, L.N.Lipatov and B.Niculescu for stimulating discussions and N.Armesto for fruitful collaboration.

6 References

1. M.A.Braun, Z.Phys. **C71** (1996) 123.
2. M.A.Braun and G.P.Vacca, Bologna univ. preprint, hep-ph/9711486.
3. L. N. Lipatov, JETP Lett. **59** (1994) 571.
4. L. D. Faddeev and G. P. Korchemsky, Phys. Lett. **B342** (1995) 311.
5. G. P. Korchemsky, Nucl. Phys. **B443** (1995) 255; **B462** (1996) 333.
6. J.Wosiek and R.A.Janik, preprint TPJU-21/96 (hep-th/9610208).
7. P.Gauron, L.Lipatov and B.Niculescu Phys.Lett. **B260** (1991) 407.
8. P.Gauron, L.Lipatov and B.Niculescu, Phys. Lett. **B304** (1993) 334; Z.Phys. **C63**(1994)253.
9. N.Armesto and M.A.Braun, Z.Phys. **C75** (1997) 709.

7 Table

Odderon energy and parameters of the trial function

N	ϵ	$c_2, c_3, \dots (c_1 = 1)$	ϵ_1
3	0.22865	-0.5036, 0.2895	0.21990
4	0.22632	-0.2791, -0.3190, 0.3609	0.20663
5	0.22627	-0.2009, -0.5052, 0.1557, 0.3779	0.17919
5*	0.22634	-0.2021, -0.5028, 0.1543, 0.3775	
6	0.22619	-0.3735, 0.08842, -0.9765, 1.003, 0.3471	-0.0015803
7	0.22618		-0.37771
8	0.22616		-0.54117
9	0.22616		-0.60273

*) double precision.